©Journal of Technical University at Plovdiv
Fundamental Sciences and Applications, Vol. 2, 1996
Series A-Pure and Applied Mathematics
Bulgaria, ISSN 1310-8271

arXiv:0808.0894v1 [math.CV] 6 Aug 2008

On some interpolation theorems for the multipliers of the Cauchy- Stiltjes type integrals

Peyo Stoilov

Abstract

Let m_α $(\alpha > 0)$ denote the set of all multipliers of the analytic functions in the unit disk D, representable by a Cauchy - Stiltjes type integral $\int\limits_T 1/(1-\overline{\zeta}z)^\alpha\, d\mu(\zeta)$.

Let the sequence $a = \{a_k\}_{k\ge 1} \subset D$ and

$$R_a f = \{f(a_k)\}_{k\ge1}, \qquad S_a f = \{f'(a_k)(1-|a_k|^2\}_{k\ge1}.$$

In this paper we prove that if a sequence $a = \{a_k\}_{k\ge1} \subset D$ satisfies conditions of Newman - Carleson and Stoltz , then

$$R_a f = b_v\,, \;\; S_a f = l^1.$$

1 Introduction

Let D denote the unit disk in the complex plane and T - the unit circle. For $0 < p \le \infty$ let H^P be the usual Hardy class [1].

Let M be the Banach space of all complex-valued Borel measures on T with the usual variation norm. For $\alpha > 0$, let F_α denote the family of all functions g for which there exists $\mu \in M$ such that

$$g(z) = \int\limits_T \frac{1}{(1-\overline{\zeta}z)^\alpha}\, d\mu(\zeta), \quad z \in D. \tag{1}$$

---

[0] **1991 Mathematics Subject Classification:** Primary 30E20, 30D50.
[0] *Key words and phrases:* Free interpolation, Cauchy - Stiltjes type integral, multipliers.

We note that F_α is a Banach space with the natural norm

$$\|g\|_{F_\alpha} = \inf\{\|\mu\| : \ \mu \in M \ \textit{such that} \ (1) \ \textit{holds}\,\}.$$

The family F_α was introduced in [2] and the following results were proved there:

If $f \in F_\alpha$, $g \in F_\beta$, then $fg \in F_{\alpha+\beta}$ $(\alpha > 0, \ \beta > 0)$.

If $\alpha < \beta$, then $F_\alpha \subset F_\beta$.

Definition. *Suppose that f is holomorphic in D . Then f is called a multiplier of F_α if $g \in F_\alpha \Rightarrow fg \in F_\alpha$.*

Let m_α denote the set of all multipliers of F_α and

$$\|f\|_{m_\alpha} = \sup\left\{\|fg\|_{F_\alpha} : \|g\|_{F_\alpha} \leq 1\right\}.$$

Various properties of m_α were studied in [3], [8]. For example it is proved that $\alpha < \beta$ implies $m_\alpha \subset m_\alpha \subset H^\infty$. Also it is proved that if $f' \in H^1$, then $f \in m_\alpha$ for all $\alpha > 0$.

These results will be applied here.

In this paper some interpolation theorems for m_1 and F_1 obtained in [4] and [7] are generalized for m_α and F_α .

2 Interpolation theorems

Definition. *We say that a sequence $a = \{a_k\}_{k\geq 1} \subset D$ satisfies Newman - Carleson condition (condition ($N - C$)) if*

$$\delta(a) = \inf\ \Big(\prod_{k\neq n} \left|\frac{a_k - a_n}{1 - \overline{a}_k a_n}\right| : \ n = 1, 2, \ldots \Big) \ > 0.$$

Definition. *We say that a sequence $a = \{a_k\}_{k\geq 1} \subset D$ satisfies the Stoltz condition (condition (S)) if*

$$\Lambda(a) = \sup\ \Big(\frac{|1 - a_k|}{1 - |a_k|} : \ k = 1, 2, \ldots \Big) \ < \infty.$$

We will need the following Lemma.

Lemma 1. *If a sequence* $a = \{a_k\}_{k\geq 1} \subset D$ *satisfy conditions* ($N - C$) *and* (S) *then*

$$\Omega(a) \; = \; \sup_{y\geq 0} \; \sum_{k\geq 1} \frac{(1-|a_k|)y}{(1-|a_k|)^2 + y^2} < \infty.$$

The proof of this Lemma can be found, for example in [5] (10.2 Lemma 11) or [6].

Let l^1 and b_v is the class of sequences:

$$l^1 = \; (x = \{x_k\}_{k\geq 1} : \quad \|x\|_{l^1} = \sum_{k\geq 1} |x_k| \; < \; \infty \;),$$

$$b_v = \; (x = \{x_k\}_{k\geq 1} : \quad \|x\|_{b_v} = \sum_{k\geq 1} |x_{k+1} - x_k| \; < \; \infty \,).$$

The following two theorems were proved in [4] for $\alpha = 1$.

Theorem 1. *Let* $\alpha > 0$ *and the sequence* $a = \{a_k\}_{k\geq 1}$ ($a_k \neq a_n$ *if* $k \neq n$ *and* $|a_k| \leq |a_{k+1}|$) *satisfy conditions* ($N - C$) *and* (S).

Then

a) If $f \in m_\alpha$, *then* $\{f(a_k)\}_{k\geq 1} \in b_v$;

b) For each sequence $x = \{x_k\}_{k\geq 1} \in b_v$, *there is a function* f *in* m_α , *such that* $f(a_k) = x_k, \quad k = 1, 2, \ldots .$

Proof. a) Let $f \in m_\alpha$. Then

$$\|\{f(a_k)\}\|_{b_v} = \sum_{k\geq 1} |f(a_k) - f(a_{k+1})| \; \leq \sum_{k\geq 1} |f(a_k) - f(|a_k|)| +$$
$$+ \sum_{k\geq 1} |f(|a_k|) - f(|a_{k+1}|)| + \sum_{k\geq 1} |f(|a_{k+1}| - f(a_{k+1})| \leq 2\Lambda(f) \; + \; I(f) \; ,$$

where

$$I(f) = \sum_{k\geq 1} |f(|a_k|) - f(|a_{k+1}|)| \, ,$$

$$\Lambda(f) \;\; = \sum_{k\geq 1} |f(a_k) - f(|a_k|)| \, .$$

In [3] (Theorem 2.6) it was proved that $\int_0^1 |f'(r)| \, dr < \infty$ and consequently

$$I(f) \leq \sum_{k\geq 1} \int\limits_{|a_{k+1}|}^{|a_k|} |f'(r)| \, dr < \int_0^1 |f'(r)| \, dr < \infty.$$

We will show that $\Lambda(f) < \infty$ too.
Let $n \geq [\alpha] + 1$ is a natural number. Then $m_\alpha \subset m_n$ and $f \in m_n$.
Since $\|1/(1-z)^n\|_{F_n} = 1$, then there exists a measurer $\mu \in M$ for which

$$\frac{f(z)}{(1-z)^n} = \int\limits_T \frac{1}{(\zeta - z)^n} d\mu(\zeta) \Rightarrow f(z) = \int\limits_T \left(\frac{1-z}{\zeta - z}\right)^n d\mu(\zeta).$$

For $\Lambda(f)$ we obtain

$$\begin{aligned}\Lambda(f) &= \sum_{k\geq 1} \left| \int\limits_T \left(\frac{1-a_k}{\zeta - a_k}\right)^n - \left(\frac{1-|a_k|}{\zeta - |a_k|}\right)^n \, d\mu(\zeta) \right| \leq \\ &\leq \sum_{k\geq 1} \int\limits_T \left| \frac{1-a_k}{\zeta - a_k} - \frac{1-|a_k|}{\zeta - |a_k|} \right| L(\zeta)\, d\,|\mu|\,(\zeta) = \\ &= \sum_{k\geq 1} \int\limits_T \frac{|1-\varsigma|\,|a_k - |a_k||}{|\zeta - a_k|\,|\zeta - |a_k||} L(\zeta)\, d\,|\mu|\,(\zeta),\end{aligned}$$

where

$$\begin{aligned}L(\zeta) &= \sum_{k=0}^{n-1} \left|\frac{1-a_k}{\zeta - a_k}\right|^k \left|\frac{1-|a_k|}{\zeta - |a_k|}\right|^{n-k-1} \leq \\ &\leq \sum_{k=0}^{n-1} \left|\frac{1-a_k}{1-|a_k|}\right|^k \leq \sum_{k=0}^{n-1} \lambda^k = \frac{1-\lambda^n}{1-\lambda}.\end{aligned}$$

Therefore

$$\Lambda(f) \leq \frac{1-\lambda^n}{1-\lambda} \sum_{k\geq 1} \int\limits_T \frac{|1-\zeta|\,|a_k - |a_k||}{|\zeta - a_k|\,|\zeta - |a_k||} \, d\,|\mu|\,(\zeta).$$

Using the condition (S), we obtain

$$\frac{|a_k - |a_k||}{1-|a_k|} = \frac{|a_k - 1 + 1 - |a_k||}{1-|a_k|} \leq 1 + \lambda, \tag{2}$$

$$\left|\frac{\zeta - |a_k|}{\zeta - a_k}\right| \leq 1 + \frac{|a_k - |a_k||}{|\zeta - a_k|} \leq 1 + \frac{|a_k - |a_k||}{1-|a_k|} \leq 2 + \lambda. \tag{3}$$

It follows from the above inequalities that

$$\Lambda(f) \;\le\; \frac{1-\lambda^n}{1-\lambda}\,(1+\lambda)(2+\lambda)\int\limits_T \sum_{k\ge1} \frac{|1-\zeta|\,(\,1-|a_k|\,)}{\left|\zeta-|a_k|\right|^2}\; d\,|\mu|\,(\zeta)\;\le$$

$$\le\; \frac{1-\lambda^n}{1-\lambda}\,(1+\lambda)(2+\lambda)\,\|\mu\|\;\sup_{\zeta\in T}\sum_{k\ge1}\frac{|1-\zeta|\,(\,1-|a_k|\,)}{\left|\zeta-|a_k|\right|^2}\le$$

$$\le \frac{1-\lambda^n}{1-\lambda}\,(1+\lambda)(2+\lambda)\,\|\mu\|\;\;\sup_{\zeta\in T}\sum_{k\ge1}\frac{|1-\zeta|\,(\,1-|a_k|\,)}{(1-|a_k|)^2+|a_1|^2\,|1-\zeta|^2}.$$

Here we used the inequalities

$$\left|\zeta-|a_k|\right|^2=(\zeta-|a_k|)(\overline{\zeta}-|a_k|)=1-|a_k|\,(\zeta+\overline{\zeta})+|a_k|^2=(1-|a_k|)^2+|a_k|\,(\,2-\zeta-\overline{\zeta}\,)=$$

$$=(1-|a_k|)^2+|a_k|\,(1-\zeta)(1-\overline{\zeta})=(1-|a_k|)^2+|a_k|\,|1-\zeta|^2\ge(1-|a_k|)^2+|a_k|^2\,|1-\zeta|^2\ge$$

$$\ge(1-|a_k|)^2+|a_1|^2\,|1-\zeta|^2\,.$$

Applying Lemma 1 for $y=|a_1|\,|1-\zeta|$, we obtain

$$\Lambda(f)\;\;\le\;\;\|f\|_{m_n}\,\frac{1-\lambda^n}{1-\lambda}\,(1+\lambda)(2+\lambda)\;\Omega(\{f(a_k)\})/\,|a_1|\quad<\infty.$$

Consequently

$$\|\{f(a_k)\}\|_{b_v}\;\;\le 2\Lambda(f)\;\;+\;\;I(f)<\infty\;\;\text{and}\;\{f(a_k)\}_{k\ge1}\in b_v\;.$$

b) Let $x=\{x_k\}_{k\ge1}\in b_v$. From the theorem of Vinogradov [6] it follows that there exists an analytic function f , such that $f'\in H^1$ and $f(a_k)=x_k,\;\; k=1,2,\dots.$ Since $f'\in H^1$, it follows that $f\in m_\alpha$ for all $\alpha>0$ [3].

Theorem 2. *Let $\alpha>0$ and the sequence $a=\{a_k\}_{k\ge1}$ ($a_k\ne a_n$ if $k\ne n$ and $|a_k|\le|a_{k+1}|$) satisfy conditions ($N-C$) and (S).*

Then

a) If $g\in F_\alpha$, then $\{(1-a_k)^\alpha\,g(a_k)\}_{k\ge1}\in b_v$;

b) For each sequence $x=\{x_k\}_{k\ge1}\in b_v$, there is a function g in F_α , such that $(1-a_k)^\alpha g(a_k)=x_k,\;\; k=1,2,\dots.$

Proof. a) Let $g \in F_\alpha$ and $f(z) = (1-z)^\alpha \, g(z)$. If $n \geq [\alpha] + 1$ is a natural number, then

$$\frac{f(z)}{(1-z)^n} = \frac{f(z)}{(\zeta - z)^\alpha} \cdot \frac{1}{(\zeta - z)^{n-\alpha}} = g(z) \cdot \frac{1}{(\zeta - z)^{n-\alpha}} \in F_{\alpha + n - \alpha} = F_n$$

and from the proof of Theorem 1 follows

$$\{f(a_k)\}_{k \geq 1} = \{(1-a_k)^\alpha \, g(a_k)\}_{k \geq 1} \in b_v.$$

b) Let $x = \{x_k\}_{k \geq 1} \in b_v$. From Theorem 1 it follows that there exists a function $f \in m_\alpha$ such that $f(a_k) = x_k, \quad k = 1, 2, ...$ Then $g(z) = \frac{f(z)}{(1-z)^\alpha} \in F_\alpha$ and $(1 - a_k)^\alpha g(a_k) = x_k, \quad k = 1, 2, \ldots.$

The following theorem generalizes a result in [7].

Theorem 3. *Let* $\alpha > 0$ *and the sequence* $a = \{a_k\}_{k \geq 1}$ *satisfy conditions ($N - C$) and (S).*
Then
a) If $f \in m_\alpha$, *then* $\{ \, f'(a_k)(1 - |a_k|^2 \, \}_{k \geq 1} \in l_1$;
b) For each sequence $x = \{x_k\}_{k \geq 1} \in l_1$, *there is a function* f *in* m_α *such that* $f'(a_k)(1 - |a_k|^2) = x_k, \quad k = 1, 2, \ldots.$

Proof. a) Let $f \in m_\alpha$ and $n \geq [\alpha] + 1$ is a natural number. Since $f \in m_n$ then there exists a measure $\mu \in M$ for which

$$\frac{f(z)}{(1-z)^n} = \int\limits_T \frac{1}{(\zeta - z)^n} d\mu(\zeta);$$

$$f(z) = \int\limits_T \left(\frac{1-z}{\zeta - z} \right)^n d\mu(\zeta);$$

$$f'(z) = n \int\limits_T \left(\frac{1-z}{\zeta - z} \right)^{n-1} \frac{1-\zeta}{(\zeta - z)^2} d\mu(\zeta).$$

Using the condition (S) and (3) , we obtain

$$\sum_{k \geq 1} |f'(a_k)| \, (1 - |a_k|^2) \leq n \int\limits_T \sum_{k \geq 1} \left| \frac{1 - a_k}{\zeta - a_k} \right|^{n-1} \frac{|1 - \zeta| \, (1 - |a_k|^2)}{|\zeta - a_k|^2} \; d \, |\mu| \, (\zeta) \;\; \leq$$

$$\leq 2n\lambda^{n-1}(2+\lambda)^2 \int\limits_T \sum_{k\geq 1} \frac{|1-\zeta|\,(1-|a_k|)}{|\zeta-|a_k||^2}\ d\,|\mu|\,(\zeta)$$

$$\leq 2n\lambda^{n-1}(2+\lambda)^2\,\|\mu\| \ \sup_{\zeta\in T} \sum_{k\geq 1} \frac{|1-\zeta|\,(\,1-|a_k|\,)}{|\zeta-|a_k||^2}$$

$$\leq 2n\lambda^{n-1}(2+\lambda)^2\,\|\mu\| \ \sup_{\zeta\in T} \sum_{k\geq 1} \frac{|1-\zeta|\,(\,1-|a_k|\,)}{(1-|a_k|)^2+|a_1|^2\,|1-\zeta|^2}.$$

Applying Lemma 1 for $y=|a_1|\,|1-\zeta|$, we obtain

$$\sum_{k\geq 1} |f'(a_k)|\,(1-|a_k|^2) \ \leq \ \|f\|_{m_n}\, 2n\lambda^{n-1}(2+\lambda)^2\,\Omega(\{f(a_k)\})/\,|a_1| \quad < \infty.$$

Consequently $\{f(a_k)\}_{k\geq 1} \in l_1$.

b) Let $x=\{x_k\}_{k\geq 1}\in l_1$. We construct the following function

$$h(z)=\sum_{n\geq 1} \frac{(1-|a_n|^2)B_n(z)}{(1-\,\overline{a}_n\,z)^2 B_n(a_n)}x_n\,, \qquad where \quad B_n(z)=\prod_{\substack{k=1\\k\neq n}}^{\infty} \frac{z-a_k}{1-\,\overline{a}_k\,z}\cdot\frac{|a_k|}{a_k}\,.$$

Since

$$\left\|\frac{1-|a_n|^2}{(1-\,\overline{a}_n\,z)^2}\right\|_{H^1} = \frac{1}{2\pi}\int\limits_T \frac{1-|a_n|^2}{|1-\,\overline{a}_n\zeta|^2}\ |d\zeta| = \frac{1}{2\pi i}\int\limits_T \frac{1-|a_n|^2}{(1-\,\overline{a}_n\,\zeta)(\zeta-\,a_n\,)}\,d\zeta \ =1,$$

$$|B_n(a_n)|\geq\delta(a)>0,$$

then

$$\|h\|_{H^1}\leq\|x\|_{l^1}/\delta(a)\ <\infty \ and \ h\in H^1.$$

Let $f(z)=\sum_{k\geq 0}\frac{\hat{h}(k)}{k+1}\,z^{k+1}$ $\quad (\,f'(z)=h(z)\,)$.

Since $f'\in H^1$ then $f\in m_\alpha$ for all $\alpha>0$ [3] and

$$f'(a_k)(1-|a_k|^2)=h(a_k)(1-|a_k|^2)\ =\ x_k,\quad k=1,2,\ldots.$$

References

[1] J. Garnett. *Bounded analytic functions.* Academic Press, New York, 1981.

[2] T. H. MacGregor. *Analytic and univalent functions with integral representations involving complex measures.* Indiana Univ. Math. J., 36(1987), 109-130.

[3] R. A. Hibschweiler and T. H. MacGregor. *Multipliers of families of Cauchy-Stieltjes transforms.* Trans. Amer. Math. Soc., 331 (1992), 377-394.

[4] S. A. Vinogradov. *Properties of multipliers of Cauchy-Stieltjes integrals and some factorization problems for analytic functions.* Amer. Math. Soc. Transl. (2) 115 (1980), 1-32.

[5] S. A. Vinogradov and V. P. Havin. *Free interpolation in H^∞ and in certain other classes of functions. I.* Zap. Nauchn. Sem. Leningrad, LOMI. 47 (1974), 15-54; English transl. in J. Soviet Math. 9 (1978).

[6] S. A. Vinogradov. *On the interpolation of power series which converge absolutely on the boundary of the circle of convergence.* Vestnik Leningrad Univ. 20 (1965), no 7, 30-44 (Russian).

[7] P. Stoilov. *Interpolation theorems for the multipliers of the Cauchy-Stieltjes type integrals.* Mathematics and Education in Mathematics, Proc. of Conference of the Union of Bulgarian Mathematicians. (1992), 10-15.

[8] D. J. Hallenbeck, T. H. MacGregor and K. Samotij. *Fractional Cauchy Transforms, Inner functions and multipliers.* Proc. London Math. Soc. (3) 72 (1996), 157-187.

Department of Mathematics
Technical University
25, Tsanko Dijstabanov,
Plovdiv, Bulgaria
e-mail: peyyyo@mail.bg